



Bottom-Up Mentalization: Bridging the Autonomic Gap in MBT through Pre-Therapeutic Arousal Regulation and Implicit Social Cognition

Zeestraten-Bartholomeus^{1,*}, and Mehrad A²

¹ Dr. Paula Zeestraten-Bartholomeus, ReAttach Academy, Berg en Terblijt, The Netherlands

² Prof. Dr. Aida Mehrad, Universitat Internacional de Catalunya, Barcelona, Spain.

***Corresponding author:** Dr. Paula Zeestraten-Bartholomeus, ReAttach Academy, Berg en Terblijt, The Netherlands,
E-mail: paula@reattachacademy.nl

Abstract

When patients experience severe emotional dysregulation and intense physical arousal, they can reach a point where treatments like Mentalization-Based Treatment (MBT) are no longer effective. This article presents ReAttach, a brief and flexible approach that serves as a practical starting point for addressing this gap. ReAttach combines the Emotional Symptom Regulation Model (ESRM) with Mapping the Individual State of Mind (M.I.S.T.) to provide targeted support tailored to the patient's current state. Features such as the Forgive and Forget Hood (FFH) help break cycles of rumination, as seen both in changes in Q-EEG brain activity and in how patients quickly move from physical distress to feeling more in control. The New Mind Creation (NMC) protocol also encourages social understanding and secure attachment, even when patients are not ready to talk about their feelings. We describe a three-phase framework that first helps calm the body before moving on to reflective mental work, aiming to reduce therapy drop-out.

Keywords: ReAttach, Mentalization-Based Treatment (MBT), Forgive and Forget Hood (FFH), Wiring Affect with ReAttach (W.A.R.A.), New Mind Creation (NMC), autonomic regulation, rumination, qualitative language analysis

Introduction: The Mentalization Threshold and the Burden of Rumination

Conditions such as Borderline Personality Disorder (BPD), trauma, and Functional Neurological Disorder (FND) are closely linked to severe emotional dysregulation and unstable physiological states (2). Top-down treatments like Mentalization-Based Treatment (MBT) can help people understand their own and others' mental states, but these approaches work best when working memory is clear and the person is within their optimal window of tolerance.

When a person's autonomic arousal rises too high, they reach a 'mentalization threshold.' After this point, their ability to reflect and think clearly shuts down, and they fall back on more basic ways of thinking, like psychic equivalence. Ongoing rumination can also overload working memory, which is a major challenge for standard psychotherapy (6). Top-down therapies often stall if the nervous system stays stuck in a state of acute stress. To address this, clinicians need a flexible, multimodal approach that helps clear mental clutter and calm arousal before starting reflective conversations.

ReAttach offers this important physiological foundation. It is a brief, transdiagnostic, and multimodal intervention that improves basic sensory processing and creates the right conditions for active learning. By tracking a patient's current state of mind, ReAttach builds a reliable neurobiological bridge, helping the body get ready to handle and process later top-down reflection (2).

Clearing the Mind: The Forgive and Forget Hood (FFH)

When a patient's working memory is filled with intrusive thoughts and sensory overload, it is hard for them to reflect deeply. The ReAttach Forgive and Forget Hood (FFH) is a practical neuropsychological exercise that helps break these rumination cycles and bring back emotional control (6).

The Neurobiological Reset

Clinical case evaluations show that the FFH protocol has an immediate effect on both body and mind. After just one self-regulation session, people reported a big drop in how burdened they felt, with average scores going from 72% to

19.08%. Quantitative EEG (Q-EEG) scans confirmed that this relief matches clear changes in brain activity. The data showed large decreases in Theta, Alpha, and Beta frequencies in the right frontal lobe, which is important for executive control and managing emotions (6). This quick return to normal brain patterns helps people handle distress through their thinking, rather than feeling overwhelmed by emotion.

Qualitative Linguistic Shifts

When patients become more stable internally, it changes how they talk about their experiences. Their language usually shifts through four stages: they begin by describing physical symptoms, then move to clearer thinking, and later talk about their experiences in more practical ways, often with a sense of control (3). At first, people who feel hyperaroused often describe their distress in strong physical terms, saying things like they have a “full head” or feel “very heavy.” After completing the FFH sequence, these physical complaints usually go away. Patients often report a new sense of mental space, using phrases like having an “empty head” or saying the uncomfortable feeling is just “gone” (3). This change in language can happen in as little as 72 hours for people without chronic conditions, but it may take several months for those with long-term issues. This pattern shows a common process of cognitive reorganization across different groups (1, 3).

Implicit Mentalization: New Mind Creation (NMC) & Sensory Integration

Traditional MBT primarily relies on explicit verbal dialogue, whereas mentalization involves both explicit and implicit processes. ReAttach focuses on the automatic, implicit side of social cognition, so patients can develop mentalizing skills without needing to report everything verbally right away.

New Mind Creation and the “Loving Observer”

Chronic toxic stress keeps the developing brain stuck in basic survival patterns and blocks the growth of healthy mental models. The New Mind Creation (NMC) framework uses specific, state-dependent tasks to break down these rigid mindsets (4). By reducing hypervigilance and strengthening automatic safety cues, the protocol prepares the nervous system for secure attachment, which is the foundation of healthy mentalization. By encouraging a “Loving Observer” perspective, NMC helps replace toxic shame with a more compassionate view of oneself (5). This approach allows patients to look at their emotional reactions from a balanced, calm place, making it easier to move from a fragmented sense of self to a more stable and adaptive identity (5).

Mapping the Autonomic State: ESRM, M.I.S.T., and Tailored Co-Regulation

One key benefit of the ReAttach approach is that it avoids a one-size-fits-all model and instead offers fully personalized, state-sensitive interventions.

ReAtt Aff Co 2(2): 178-180 (2026)

The Emotional Symptom Regulation Model (ESRM) & M.I.S.T.

Recent digital phenotyping studies involving more than 1,000 people have identified seven main emotional clusters: Distress, Shutdown, Activation, Anger, Social Isolation, Resilience, and Ambivalence (2). This system goes beyond just distress and resilience. ReAttach uses the M.I.S.T. (Mapping the Individual State of Mind) interview to find specific processing bottlenecks (2).

The Hyper-Aroused (Distress/Anger): Patients stuck in high sympathetic activation need quick ways to calm down. Techniques like Wiring Affect with ReAttach (W.A.R.A.) help separate strong negative feelings from interpersonal triggers, and the FFH helps quiet overactive frontal brain activity (6).

The Hypo-Aroused (Shutdown/Social Isolation): People in dorsal vagal immobilization often show emotional flatness and severe brain fog. They need gentle upregulation and specific tactile rhythms to boost oxytocin levels, helping to rebuild social motivation before moving on to more challenging cognitive tasks (2).

Systemic Co-Regulation and the Safety Override Technique (SOT)

Meaningful recovery in complex situations rarely happens alone; it depends on relationships (1). ReAttach supports this by training partners or family caregivers to help with stabilization. For people experiencing sudden sensory overload or hypervigilance, the new Safety Override Technique (SOT) provides a helpful emergency protocol (1). SOT uses quick tapping and structured sensory-motor cues to interrupt threat responses and guide the nervous system back to a safer state. Since it can be practiced and co-regulated at home, it is a practical tool for daily long-term management (1).

The Proposed Clinical Integration: The 3-Phase Protocol

The researchers suggest a step-by-step, three-phase clinical workflow to help top-down therapies work more effectively and reduce patient dropout.

1. Phase 1: Autonomic Reset & Cognitive Clearing (W.A.R.A., SOT & FFH)
2. This phase helps the patient's nervous system reach a manageable state, guided by their M.I.S.T. profile. W.A.R.A. and SOT are used to calm strong emotions, and FFH addresses ongoing rumination to clear working memory (2, 6).
3. Phase 2: Implicit Mentalization (New Mind Creation)
4. This phase builds early attachment security using different types of sensory input. Creating the “Loving Observer” helps lessen the impact of unhelpful thought patterns. ReAttach has been shown to reduce these patterns, with a large effect

size ($d=1.28$), and to support a resilient growth mindset (4, 5).

5. Phase 3: Explicit MBT
6. After the patient's body is stable, working memory is clear, and their ability to track thoughts is ready, they can start the explicit, verbal Socratic work that is central to traditional MBT.

Conclusion

Patients with complex personality issues or functional disorders can be treated. In many cases, the main challenge is that their physical state does not match what traditional therapies require. ReAttach offers a practical and wide-ranging method. It maps autonomic states (ESRM/M.I.S.T.), uses systemic co-regulation, interrupts repetitive thinking (FFH), and helps build secure attachment (NMC). These steps support the patient's ability to learn (2, 4, 6). We invite clinicians everywhere to try this three-phase protocol, where bottom-up regulation and top-down mentalization come together to open new paths to recovery.

References

1. Call, E., Painter, S., Zeestraten-Bartholomeus, P., & Mehrad, A. (2026). From hopelessness to self- and co-regulation: A mixed-methods exploratory case series of remote ReAttach

in Functional Neurological Disorder. *ReAttach Affect Coach Journal* (manuscript in preparation).

2. Painter, S., Zeestraten-Bartholomeus, P., Bitá, M., & Mehrad, A. (2026). Blended insights from the ReAttach M.I.S.T. and Emotional Symptom Regulation Model (ESRM). *ReAttach Affect Coach Journal*, 2(1), 102-115.
3. Zeestraten-Bartholomeus, P., Painter, S., Lindenberg, G., & Mehrad, A. (2026). From bodily sensations to metacognitive reflection: A qualitative language analysis of ReAttach interventions. *ReAttach Affect Coach Journal* (manuscript in preparation).
4. Zeestraten-Bartholomeus, P., Painter, S., Bitá, M., & Mehrad, A. (2026). Stimulating a growth mindset and adaptive schemas in children via New Mind Creation. *ReAttach Affect Coach Journal*, 2(1), 126-129.
5. Zeestraten-Bartholomeus, P., Painter, S., Bitá, M., & Mehrad, A. (2026). From autonomic fixation to cohesive identity: Integrating the ESRM model and ReAttach therapy in the treatment of personality disorders. *ReAttach Affect Coach Journal*, 2(1), 141-144.
6. Zeestraten-Bartholomeus, P., Lee, J., Tan, D., Bitá, M., & Mehrad, A. (2025). ReAttach Forgive and Forget Hood: Clear the mind, stop rumination! *ReAttach Affect Coach Journal*, 1(2), 40-47.



ReAttach Affect Coach Journal is a peer-reviewed open access journal. © 2025, this is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.